

PACKAGE INSERT  
PLEASE SAVE FOR FUTURE USE

CONTAINS ACETAMINOPHEN + METHOCARBAMOL

# Robaxacet<sup>®</sup>

## EXTRA STRENGTH

---

**Indications:** For effective relief of pain associated with muscle spasm such as back pain, tense neck muscles, strains and sprains.

**Each caplet contains the following active ingredients:** Methocarbamol (muscle relaxant) 400 mg, **Acetaminophen** (analgesic, extra strength) 500 mg.

**DIRECTIONS: Adult Dosage (12 years and older):** 2 caplets, every 6 hours. Do not exceed 8 caplets in 24 hours. Use only on the advice of a doctor.

● **Caution: KEEP OUT OF REACH OF CHILDREN.** This package contains enough drug to seriously harm a child. **DO NOT USE** with other products containing **acetaminophen or methocarbamol**. If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. **ADULT USE ONLY.** Do not take more than the recommended dose unless advised by a doctor. Use the smallest effective dose. **Liver warning:** This product contains acetaminophen. Maximum daily dose is 8 caplets in 24 hours. **Severe or possibly fatal liver damage** may occur if you take: more than the recommended dose in 24 hours; with other drugs containing acetaminophen; while drinking three (3) or more alcoholic drinks every day. Symptoms of liver damage may include: yellowing of the skin/eyes, dark urine; sweating, nausea, vomiting, stomach pain; unusual tiredness, and/or loss of appetite. **Allergy alert:** Acetaminophen may cause serious skin reactions. Symptoms may include skin reddening, blisters and rash. **If any of the above noted symptoms occur, stop use and seek medical help right away.** Stop use and consult a doctor if pain lasts more than 5 days. **Ask a doctor before use** if you suffer from chronic alcoholism, serious liver or kidney disease. **Ask a doctor or pharmacist before use:** if you take Warfarin-containing blood thinning drugs, if pregnant, breast feeding or under a physician's care. Do not use with a monamine oxidase inhibitor (MAOI) or for two weeks after stopping the MAOI drug. Consult a doctor prior to combining with medication for high blood pressure, depression or any other medication including prescription drugs, salicylates, nasal decongestants or pain medications. May cause drowsiness or dizziness or nausea. Consult a doctor before use if taking sleeping pills, sedatives or tranquilizers, or cold and

allergy medications, since these medications may increase drowsiness. Avoid alcoholic beverages, driving a motor vehicle or operating machinery.

**In Case of Overdose:** Call a Poison Control Centre or get medical help right away. Quick medical attention is critical even if you do not notice any signs or symptoms.

**Non-Medicinal Ingredients:** Corn starch, crospovidone, D&C Yellow No. 10, FD&C Blue No. 1, hypromellose, magnesium stearate, methylcellulose, polyethylene glycol, povidone, pregelatinized starch, silicon dioxide, sodium lauryl sulfate, sodium starch glycolate, stearic acid.

Store at room temperature (15°-30°C).

Pfizer Consumer Healthcare, a division of Pfizer Canada Inc.,  
Mississauga, ON, Canada L5R 4B2

Questions? 1-888-275-9938 [www.backrelief.ca](http://www.backrelief.ca)