

Robaxacet®

Acetaminophen + Methocarbamol

This is the most current labeling information and may differ from labels on product packaging. If there are any differences between this website labeling and product packaging labeling, this website labeling should be regarded as the most current.

Drug Facts

Active ingredients (in each caplet)	Purposes
Acetaminophen (APAP) 325 mg	Pain reliever
Methocarbamol 400 mg.....	Muscle relaxant

Uses •For temporary relief of pain associated with muscle spasm such as
■back pain ■tense neck muscles ■strains and sprains.

Warnings **Liver warning** Acetaminophen may cause **severe or possibly fatal liver damage** if you take: •more than the recommended dose in 24 hours •with other drugs containing acetaminophen •while drinking 3 or more alcoholic drinks every day. Symptoms of liver damage may include: •yellow skin or eyes •dark urine •sweating •nausea •vomiting •stomach pain •unusual tiredness •loss of appetite.

Allergy alert Acetaminophen may cause serious skin reactions. Stop use and get medical help right away if you have: •skin reddening •blisters •rash. **If any of the above noted symptoms occur, stop use and seek medical help right away.**

Do not use •with any other drugs containing acetaminophen or other ingredients in this product •if allergic to acetaminophen or other ingredients in this product. If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

Ask a doctor or pharmacist before use if you •have liver or kidney disease •take ■ the blood thinning drug warfarin ■drugs for depression, sedatives or any other drug •are pregnant or breastfeeding.

When using this product •drowsiness, dizziness, nausea or light-headedness may occur •do not drive or operate machinery •do not drink alcohol.

Stop use and ask a doctor if •symptoms worsen or pain lasts for more than 5 days •or get any unusual symptoms •you get a rash or itchy skin.

Keep out of reach of children. This package contains enough drug to seriously harm a child. In case of overdose, call a poison control centre or get medical help right away. Quick medical attention is critical even if you do not notice any signs or symptoms.

Directions Adults and children 12 years & over: •take 2 caplets every 6 hours •do not take more than 8 caplets in 24 hours. Do not take more than directed (see liver warnings). Use only on the advice of a doctor.

Other information Store at room temperature (15°-30°C).

Inactive ingredients Corn starch, crospovidone, D&C Yellow No. 10, FD&C Blue No. 1, hypromellose, magnesium stearate, methylcellulose, polyethylene glycol, povidone, pregelatinized starch, silicon dioxide, sodium lauryl sulfate, sodium starch glycolate, stearic acid.

Questions? 1-888-275-9938 www.robax.ca